

ALMOST AUTOMORPHIC SOLUTIONS TO ABSTRACT VOLTERRA EQUATIONS ON THE LINE

CARLOS LIZAMA AND RODRIGO PONCE

ABSTRACT. Given $a \in L^1(\mathbb{R})$ and A the generator of an L^1 -integrable family of bounded and linear operators defined on a Banach space X , we prove the existence of almost automorphic mild solution to the semilinear integral equation $u(t) = \int_{-\infty}^t a(t-s)[Au(s) + f(s, u(s))]ds$ for each $f : \mathbb{R} \times X \rightarrow X$ S^p -almost automorphic in t , uniformly in $x \in X$, and satisfying diverse Lipschitz type conditions. In the scalar linear case, we prove that $a \in L^1(\mathbb{R})$ completely monotonic is already sufficient.

1. INTRODUCTION

We study almost automorphic solutions of an integral equation with infinite delay in a general Banach space X :

$$(1.1) \quad u(t) = \int_{-\infty}^t a(t-s)[Au(s) + f(s, u(s))]ds, \quad t \in \mathbb{R}$$

where the operator $A : D(A) \subset X \rightarrow X$ generates an integral resolvent and $a : \mathbb{R}_+ \rightarrow \mathbb{C}$ is an integrable function.

A rich source of problems leading to the equation (1.1) is provided by the theory of viscoelastic material behavior. Some typical examples are viscoelastic fluids and heat flow in materials of fading memory type: see for instance [3], [24] and [27]. In such applications the operator A typically is the Laplacian in $X = L^2(\Omega)$, or the elasticity operator, the Stokes operator, or the biharmonic Δ^2 , etc. equipped with suitable boundary conditions. The material kernel $a(t)$ reflect the properties of the medium under consideration. Note that, in the finite dimensional case, the system (1.1) contains as particular cases several systems with finite or infinite delay, already considered in the literature. See e.g. [4] and [16].

An equivalent form of equation (1.1) is given by

$$(1.2) \quad u(t) + \frac{d}{dt}(\alpha u(t) + \int_{-\infty}^t k(t-s)u(s)ds) = \int_0^\infty a(s)ds(Au(t) + f(t, u(t))), \quad t \in \mathbb{R}$$

for some $\alpha > 0$ and $k \in L^1(\mathbb{R}_+)$ nonnegative and non increasing, see [5, Section 2]. This integrodifferential equation was studied in [6], where some results of [7] were used in order to obtain the existence and regularity of the solution u when A generates a contraction semigroup (not necessarily analytic) on X .

2000 *Mathematics Subject Classification.* Primary 34G20; Secondary 45N05, 43A60.

Key words and phrases. S^p -almost automorphic function; integral resolvent family; semilinear integrodifferential equations.

The first author is partially supported by FONDECYT under Grant number: 1100485.

The problem of existence of almost automorphic solutions to (1.1) is a very natural one. Conditions which guarantee the existence of an almost automorphic solution for any $f(t, x)$ in a given space of almost automorphic functions have been studied recently in [8].

In a recent paper [13], the authors dealt with the existence of almost automorphic solutions to certain classes of fractional differential equations, which can be represented in the form ([9, Section 1]):

$$u(t) = \int_0^t \frac{(t-s)^{\alpha-1}}{\Gamma(\alpha)} [Au(s) + g(s, u(s))] ds; \quad u(0) = u_0, \quad 1 < \alpha < 2.$$

The aim of this paper is to point out that similar results hold true for the class of integral equations (1.1) (or equivalently (1.2)) containing the above equations as limiting special cases [27, Chapter II, section 11.5].

Specifically, we consider in this paper the class of continuous data $f : \mathbb{R} \times X \rightarrow X$ of S^p -almost automorphic functions on t , and we look for solutions u belonging to the class of almost automorphic functions.

The concept of S^p -almost automorphy was introduced and applied to study the existence of solutions to some parabolic evolution equations by N'Guérékata and Pankov in [23]. We would like to point out that new and interesting results on S^p -almost automorphic mild solutions to evolution equations have been recently appeared in [17], [11] and [18]. However, none of them include existence of almost automorphic mild solutions for (1.1) or (1.2) with S^p -almost automorphic terms.

This paper is organized as follows: Section 2 collect all the results of [25],[11], [23], [13] and [12] we need about S^p -almost automorphic functions. In section 3, we treat equation (1.1) when $f(t, u(t)) = g(t)$ is S^p -almost automorphic, that is, the linear case. We exploit in full strength the use of integral resolvents, instead the classical approach by C_0 -semigroups, to obtain a representation of the solution. Then, using recent results from [13] (see Lemma 2.9 below) we get maximal regularity of the solution (Proposition 3.1). In particular, we improve in this section some results of [8] and give new results and examples, specially in the scalar case, where the assumptions given are based only in the data of the equation. Specifically we require only $a \in L^1(\mathbb{R})$ completely monotonic (Theorem 3.2). Section 4 is devoted to our main results in the semilinear case. As is well known, maximal regularity results for the linear case, are very useful for nonlinear problems. Using this approach, and a very recent composition theorem proved in [13] (Theorem 2.10), we are able to prove three new and general existence and uniqueness theorems of almost automorphic solutions to the equation (1.1) (cf. Theorems 4.2, 4.3 and 4.4). Then, as an illustration, we apply the abstract results to a concrete equation. Finally, we point out that our results also generalizes the existence results obtained in [8], as the space of S^p -almost automorphic functions contains the space $AA(X)$ of almost automorphic functions.

2. PRELIMINARIES

Since we are going to use several concepts of almost automorphy, it is proper to provide a brief description of the spaces of almost automorphic functions to be considered.

Definition 2.1. A continuous function $f : \mathbb{R} \rightarrow X$ is said to be almost automorphic if for every sequence of real numbers $(s'_n)_{n \in \mathbb{N}}$ there exists a subsequence $(s_n)_{n \in \mathbb{N}} \subset (s'_n)_{n \in \mathbb{N}}$

such that

$$g(t) := \lim_{n \rightarrow \infty} f(t + s_n)$$

is well defined for each $t \in \mathbb{R}$, and

$$f(t) = \lim_{n \rightarrow \infty} g(t - s_n), \quad \text{for each } t \in \mathbb{R}.$$

If the convergence above is uniform in $t \in \mathbb{R}$, then f is almost periodic in the classical Bochner's sense.

Almost automorphy, as a generalization of the classical concept of an almost periodic function, was introduced in the literature by S. Bochner and recently studied by several authors, including [1, 2, 10, 14, 19, 20] among others. A complete description of their properties and further applications to evolution equations can be found in the monographs [21] and [22] by G. M. N'Guérékata.

Denote by $AA(X)$ the collection of all (Bochner) almost automorphic functions $\mathbb{R} \rightarrow X$. Some basic properties are stated in the following result.

Theorem 2.2 ([22]). *If $f, f_1, f_2 \in AA(X)$, then:*

- (i) $f_1 + f_2 \in AA(X)$;
- (ii) $\lambda f \in AA(X)$ for any scalar λ ;
- (iii) $f_\alpha \in AA(X)$ where $f_\alpha : \mathbb{R} \rightarrow X$ is defined by $f_\alpha(\cdot) = f(\cdot + \alpha)$;
- (iv) the range $R_f := \{f(t) : t \in \mathbb{R}\}$ is relatively compact in X , thus f is bounded in norm;
- (v) if $f_n \rightarrow f$ uniformly on \mathbb{R} where each $f_n \in AA(X)$, then $f \in AA(X)$ too.

Note that $AA(X)$ equipped with the sup-norm $\|f\|_{AA(X)} = \sup_{t \in \mathbb{R}} \|f(t)\|$ turns out to be a Banach space.

Definition 2.3 ([25]). The Bochner transform $f^b(t, s), t \in \mathbb{R}, s \in [0, 1]$, of a function $f(t)$ on \mathbb{R} , with values in X , is defined by

$$f^b(t, s) := f(t + s).$$

Definition 2.4 ([25]). The space $BS^p(X)$ of all Stepanov bounded functions, with the exponent p , consists of all measurable functions $f : \mathbb{R} \rightarrow X$ such that

$$\|f\|_{S^p} := \sup_{t \in \mathbb{R}} \left(\int_t^{t+1} \|f(\tau)\|^p d\tau \right)^{\frac{1}{p}} < \infty.$$

It is obvious that $L^p(\mathbb{R}; X) \subset BS^p(X) \subset L^p_{loc}(\mathbb{R}; X)$ and $BS^p(X) \subset BS^q(X)$ whenever $p \geq q \geq 1$.

Definition 2.5 ([23]). The space $AS^p(X)$ of S^p -almost automorphic functions (S^p -a.a. for short) consists of all $f \in BS^p(X)$ such that $f^b \in AA(L^p([0, 1]; X))$. In other words, a function $f \in L^p_{loc}(\mathbb{R}; X)$ is said to be S^p -almost automorphic if its Bochner transform $f^b : \mathbb{R} \rightarrow L^p([0, 1]; X)$ is almost automorphic in the sense that for every sequence of real numbers $(s'_n)_{n \in \mathbb{N}}$, there exist a subsequence $(s_n)_{n \in \mathbb{N}} \subset (s'_n)_{n \in \mathbb{N}}$ and a function $g \in L^p_{loc}(\mathbb{R}; X)$ such that

$$\lim_{n \rightarrow \infty} \left(\int_0^1 \|f(t + s_n + s) - g(t + s)\|^p ds \right)^{\frac{1}{p}} = 0,$$

$$\lim_{n \rightarrow \infty} \left(\int_0^1 \|g(t - s_n + s) - f(t + s)\|^p ds \right)^{\frac{1}{p}} = 0,$$

for each $t \in \mathbb{R}$.

Remark 2.6. It is clear that if $1 \leq p < q < \infty$ and $f \in L_{loc}^q(\mathbb{R}; X)$ is S^q -almost automorphic, then f is S^p -almost automorphic. Also if $f \in AA(X)$, then f is S^p -almost automorphic for any $1 \leq p < \infty$.

Denote $l^\infty(X)$ the space of all bounded sequences with values in X . Recall that a sequence $x \in l^\infty(X)$ is said to be almost automorphic if for any sequence of integers (s'_n) there exists a subsequence (s_n) such that $\lim_{n \rightarrow \infty} \lim_{m \rightarrow \infty} x_{p-s_n-s_m} = x_p$.

Example 2.7 ([23]). Let (a_n) be an almost automorphic sequence and $\epsilon_0 \in (0, 1/2)$. Let $f(t) = a_n$ if $t \in (n - \epsilon_0, n + \epsilon_0)$ and $f(t) = 0$ otherwise. Then $f \in AS^q(X)$ for all $q \in [1, \infty)$ but f is not in $AA(X)$.

Definition 2.8 ([11]). A function $f : \mathbb{R} \times X \rightarrow X$, $(t, u) \mapsto f(t, u)$ with $f(\cdot, u) \in L_{loc}^p(\mathbb{R}, X)$ for each $u \in X$ is said to be S^p -almost automorphic in $t \in \mathbb{R}$ uniformly for $u \in X$, if for every sequence of real numbers $(s'_n)_{n \in \mathbb{N}}$, there exist a subsequence $(s_n)_{n \in \mathbb{N}} \subset (s'_n)_{n \in \mathbb{N}}$ and a function $g : \mathbb{R} \times X \rightarrow X$ with $g(\cdot, u) \in L_{loc}^p(\mathbb{R}, X)$ such that

$$\lim_{n \rightarrow \infty} \left(\int_0^1 \|f(t + s_n + s, u) - g(t + s, u)\|^p ds \right)^{\frac{1}{p}} = 0,$$

$$\lim_{n \rightarrow \infty} \left(\int_0^1 \|g(t - s_n + s, u) - f(t + s, u)\|^p ds \right)^{\frac{1}{p}} = 0,$$

for each $t \in \mathbb{R}$ and for each $u \in X$. We denote by $AS^p(\mathbb{R} \times X, X)$ the set of all such functions.

Lemma 2.9 ([13]). Let $\{S(t)\}_{t \geq 0} \subset \mathcal{B}(X)$ be a strongly continuous family of bounded and linear operators such that

$$\|S(t)\| \leq \phi(t), \quad \text{for all } t \in \mathbb{R}_+,$$

where $\phi \in L^1(\mathbb{R}_+)$ is nonincreasing. Then, for each $f \in AS^1(X)$,

$$\int_{-\infty}^t S(t-s)f(s)ds \in AA(X).$$

Theorem 2.10 ([13, 12]). Assume that

(i) $f \in AS^p(\mathbb{R} \times X, X)$ with $p > 1$;

(ii) there exists a non negative function $L \in AS^r(\mathbb{R})$ with $r \geq \max\{p, p/(p-1)\}$ such that for all $u, v \in X$ and $t \in \mathbb{R}$,

$$\|f(t, u) - f(t, v)\| \leq L(t)\|u - v\|;$$

(iii) $x \in AS^p(X)$ and $K = \overline{\{x(t) : t \in \mathbb{R}\}}$ is compact in X .

Then there exists $q \in [1, p)$ such that $f(\cdot, x(\cdot)) \in AS^q(X)$.

The following definition is taken from [27, Definition 1.6, p.46].

Definition 2.11. Let X be a complex Banach space, A a closed linear unbounded operator in X and $a \in L^1_{loc}(\mathbb{R}_+)$ an scalar kernel $\not\equiv 0$. A family $\{S(t)\}_{t \geq 0} \subset \mathcal{B}(X)$ is called an integral resolvent with generator A if the following conditions are satisfied.

- (i) $S(\cdot)x \in L^1_{loc}(\mathbb{R}_+; X)$ for each $x \in X$ and $\|S(t)\| \leq \psi(t)$ a.e. on \mathbb{R}_+ , for some $\psi \in L^1_{loc}(\mathbb{R}_+)$;
- (ii) $S(t)$ commutes with A for each $t \geq 0$;
- (iii) the following integral resolvent equation holds

$$(2.1) \quad S(t)x = a(t)x + \int_0^t a(t-s)AS(s)x ds.$$

for all $x \in D(A)$ and a.a. $t \geq 0$.

We will see that the concept of integral resolvent is directly and naturally related with the solution of the equation (1.1) by means of a kind of variation of parameters formula (cf. Definition 4.1 below). On the other side, our definition of solution (mild) for equation (1.1) that we will give, is motivated by the linear case (see (3.3) in Proposition 3.1 below).

3. ALMOST AUTOMORPHIC SOLUTIONS FOR THE LINEAR EQUATION

In this section we consider the existence and uniqueness of almost automorphic solutions to the evolution equation

$$(3.1) \quad u(t) = \int_{-\infty}^t a(t-s)[Au(s) + g(s)]ds, \quad t \in \mathbb{R},$$

where A is the generator of an integral resolvent family and $a \in L^1(\mathbb{R})$.

Proposition 3.1. *Let $a \in L^1(\mathbb{R})$. Assume that A generates an integral resolvent family $\{S(t)\}_{t \geq 0}$ on X , which satisfies*

$$\|S(t)\| \leq \phi(t), \quad \text{for all } t \in \mathbb{R}_+,$$

where $\phi \in L^1(\mathbb{R}_+)$ is nonincreasing. If $f \in AS^1(X)$ and takes values on $D(A)$ then the unique bounded solution of the problem

$$(3.2) \quad u(t) = \int_{-\infty}^t a(t-s)[Au(s) + f(s)]ds, \quad t \in \mathbb{R},$$

is almost automorphic and is given by

$$(3.3) \quad u(t) = \int_{-\infty}^t S(t-s)f(s)ds, \quad t \in \mathbb{R}.$$

Proof. Since $g(t) \in D(A)$ for all $t \in \mathbb{R}$, we obtain $u(t) \in D(A)$ for all $t \in \mathbb{R}$ (see [27, Proposition 1.2]). Then applying (2.1) and Fubini's theorem we obtain

$$\begin{aligned}
\int_{-\infty}^t a(t-s)Au(s)ds &= \int_{-\infty}^t a(t-s)A \int_{-\infty}^s S(s-\tau)f(\tau)d\tau ds \\
&= \int_{-\infty}^t \int_{-\infty}^s a(t-s)AS(s-\tau)f(\tau)d\tau ds \\
&= \int_{-\infty}^t \int_{\tau}^t a(t-s)AS(s-\tau)f(\tau)ds d\tau \\
&= \int_{-\infty}^t \int_0^{t-\tau} a(t-\tau-p)AS(p)dp f(\tau)d\tau \\
&= \int_{-\infty}^t (S(t-\tau)f(\tau) - a(t-\tau)f(\tau))d\tau \\
&= u(t) - \int_{-\infty}^t a(t-\tau)f(\tau)d\tau.
\end{aligned}$$

The statement follows by Lemma 2.9. \square

Recall that a C^∞ -function $f : (0, \infty) \rightarrow \mathbb{R}$ is called completely monotonic if

$$(-1)^n f^{(n)}(\lambda) \geq 0, \text{ for all } \lambda > 0, n \in \mathbb{N}_0.$$

We remark that such functions naturally occur in areas such as probability theory, numerical analysis, and elasticity. Our main result in case $X = \mathbb{R}$ is the following theorem. It is remarkable that the hypothesis given, are completely based in the data of the problem.

Theorem 3.2. *Let $a \in L^1(\mathbb{R}_+)$ be a scalar, completely monotonic function on \mathbb{R}_+ . Let $\rho > 0$ be given. If $g \in AS^1(\mathbb{R})$ then:*

a) *There is $S_\rho \in L^1(\mathbb{R}_+)$ completely monotonic such that equation (2.1) is satisfied with $A = -\rho$.*

b) *The equation*

$$(3.4) \quad u(t) = \int_{-\infty}^t a(t-s)[- \rho u(s) + g(s)]ds, \quad t \in \mathbb{R},$$

has an unique almost automorphic solution given by

$$u(t) = \int_{-\infty}^t S_\rho(t-s)g(s)ds, \quad t \in \mathbb{R}.$$

Proof. By the hypothesis on the scalar kernel $a(t)$ and [16, Theorem 2.8, p.147] we have that $\log(a)$ is convex on \mathbb{R}_+ . Moreover, since $a(t)$ is positive and nonincreasing, it follows by [27, Lemma 4.1, p.98] that there exists $S_\rho \in L^1(\mathbb{R}_+)$ completely monotone, such that equation (2.1) is satisfied with $A = -\rho$, that is

$$(3.5) \quad S_\rho(t) = a(t) - \rho \int_0^t a(t-s)S_\rho(s)ds.$$

Hence (a) follows. Part (b) is an immediate consequence of Lemma 2.9, since S_ρ is non-increasing. \square

In case that $g \in AA(\mathbb{R})$ we have the following result that improves [8, Corollary 3.7]. We denote by $\hat{a}(\lambda)$ the Laplace transform of $a(t)$.

Theorem 3.3. *Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be an almost automorphic function and let $\rho > 0$ be a real number. Suppose $a \in L^1(\mathbb{R}_+)$, and $\hat{a}(\lambda) \neq -\frac{1}{\rho}$ for all $\operatorname{Re}(\lambda) \geq 0$. Then*

- a) *There is $S_\rho \in L^1(\mathbb{R}_+)$ such that equation (2.1) is satisfied with $A = -\rho$;*
- b) *The equation*

$$(3.6) \quad u(t) = \int_{-\infty}^t a(t-s)[- \rho u(s) + f(s)]ds, \quad t \in \mathbb{R},$$

has an unique almost automorphic solution given by

$$u(t) = \int_{-\infty}^t S_\rho(t-s)f(s)ds, \quad t \in \mathbb{R}.$$

Proof. The proof is a direct consequence of the half Paley-Wiener theorem [16, Theorem 4.1 p.45] and [8, Lemma 3.1] (see also the references therein). \square

Example 3.4. *Let $a(t) = e^{-bt}$, $b > 0$ and $\rho > 0$. Then $a(t)$ is completely monotonic and $\hat{a}(\lambda) = \frac{1}{\lambda+b} \neq -\frac{1}{\rho}$ for all $\operatorname{Re}(\lambda) \geq 0$. Moreover a direct calculation using Laplace transform gives $S_\rho(t) = e^{-(b+\rho)t}$. Hence for any $g \in AS^1(\mathbb{R})$ (resp. $g \in AA(\mathbb{R})$) there exist a unique almost automorphic solution of the equation*

$$u(t) = \int_{-\infty}^t e^{-b(t-s)}[-\rho u(s) + g(s)]ds, \quad t \in \mathbb{R},$$

given by

$$u(t) = \int_{-\infty}^t e^{(t-s)(b+\rho)}g(s)ds, \quad t \in \mathbb{R}.$$

The remarkable fact is that we only need $g \in AS^1(\mathbb{R})$ instead of $g \in AA(\mathbb{R})$ to have existence of almost automorphic solutions.

Example 3.5. *Let $a(t) = \frac{t^{\alpha-1}}{\Gamma(\alpha)}e^{-bt}$, $b > 0$, $\alpha > 0$ and $\rho > 0$. We note that $a(t)$ is not completely monotonic but, under the condition $\cos(\pi/\alpha) \leq \frac{b}{\rho^{1/\alpha}}$ and since $(-\rho)^{\frac{1}{\alpha}} = [\cos(\frac{\pi}{\alpha}) + i \sin(\frac{\pi}{\alpha})]\rho^{\frac{1}{\alpha}}$, we have $\lambda \neq (-\rho)^{\frac{1}{\alpha}} - b$, for all $\operatorname{Re}(\lambda) \geq 0$, that is $\hat{a}(\lambda) = \frac{1}{(\lambda+b)^\alpha} \neq -\frac{1}{\rho}$, for all $\operatorname{Re}(\lambda) \geq 0$. A calculation using Laplace transform shows $S_\rho(t) = t^{\alpha-1}e^{-bt}E_{\alpha,\alpha}(\rho t^\alpha)$, where $E_{\alpha,\alpha}$ denotes the generalized Mittag-Leffler function (see e.g. [15]).*

Hence for any $g \in AA(\mathbb{R})$ there exist a unique almost automorphic solution of the equation

$$u(t) = \frac{1}{\Gamma(\alpha)} \int_{-\infty}^t (t-s)^{\alpha-1}e^{-b(t-s)}[-\rho u(s) + g(s)]ds, \quad t \in \mathbb{R},$$

given by

$$u(t) = \int_{-\infty}^t (t-s)^{\alpha-1}e^{-b(t-s)}E_{\alpha,\alpha}(\rho(t-s)^\alpha)g(s)ds, \quad t \in \mathbb{R}.$$

Note that this example improves [8, Example 3.6] where only $1 < \alpha < 2$ was considered. In fact, $1 < \alpha < 2$ implies immediately that the more general condition $\cos(\pi/\alpha) \leq \frac{b}{\rho^{1/\alpha}}$ holds.

Example 3.6. Let $b > 0$ and consider the perturbed problem

$$(3.7) \quad u(t) = \int_{-\infty}^t e^{-b(t-s)} [Bu(s) + bu(s) + f(s)] ds, \quad t \in \mathbb{R},$$

where B is the generator of an exponentially stable C_0 -semigroup $T(t)$. Taking $a(t) = e^{-bt}$ and $A := B + bI$ we obtain (3.7) in the form of equation (3.2). By (2.1) we have

$$\hat{S}(\lambda) = \left(\frac{1}{\hat{a}(\lambda)} - A \right)^{-1} = (\lambda + b - B - b)^{-1} = (\lambda - B)^{-1} = \hat{T}(\lambda).$$

Hence, by uniqueness of the Laplace transform, we obtain in this case that the integral resolvent $S(t)$ is identical to the C_0 -semigroup $T(t)$. We conclude by Proposition 3.1 that given $f \in AS^1(X)$ taking values on $D(B)$, the unique bounded solution of the problem (3.7) is almost automorphic and given by

$$u(t) = \int_{-\infty}^t T(t-s)f(s)ds, \quad t \in \mathbb{R}.$$

4. ALMOST AUTOMORPHIC MILD SOLUTIONS FOR NONLINEAR EQUATIONS

In this section we consider the existence and uniqueness of almost automorphic mild solutions to the nonlinear evolution equation

$$(4.1) \quad u(t) = \int_{-\infty}^t a(t-s)[Au(s) + f(s, u(s))]ds, \quad t \in \mathbb{R},$$

where A is the generator of an integral resolvent family and $a \in L^1(\mathbb{R})$. The following definition is motivated by the linear case.

Definition 4.1. Let A be the generator of an integral resolvent family $\{S(t)\}_{t \geq 0}$. A continuous function $u : \mathbb{R} \rightarrow X$ satisfying the integral equation

$$(4.2) \quad u(t) = \int_{-\infty}^t S(t-s)f(s, u(s))ds, \quad \text{for all } t \in \mathbb{R},$$

is called a *mild* solution on \mathbb{R} to the equation (4.1).

Our first result gives conditions under which we have existence of a unique almost automorphic mild solution with S^p -almost automorphic terms. Note that we assume that the function f is bounded by a Lipschitz function $L(t)$ with respect to the first argument uniformly in the second argument. Moreover, we have to assume a control on the S^1 -norm of the Lipschitz function. The advantage is that the assumptions on L and ϕ are less restrictive than in the subsequent results.

Theorem 4.2. Assume that A generates an integral resolvent family $\{S(t)\}_{t \geq 0}$ such that

$$\|S(t)\| \leq \phi(t), \quad \text{for all } t \geq 0,$$

where $\phi \in L^1(\mathbb{R}_+)$ is nonincreasing. Suppose that

- (i) $f \in AS^p(\mathbb{R} \times X, X)$ with $p > 1$;
- (ii) there exists a non negative function $L \in AS^r(\mathbb{R})$ with $r \geq \max\{p, p/(p-1)\}$ such that for all $u, v \in X$ and $t \in \mathbb{R}$,

$$\|f(t, u) - f(t, v)\| \leq L(t)\|u - v\|;$$

If $\|L\|_{S^1} < \|\phi\|_1^{-1}$, then the equation (4.1) has a unique almost automorphic mild solution.

Proof. We define the operator $F : AA(X) \mapsto AA(X)$ by

$$(4.3) \quad (F\varphi)(t) := \int_{-\infty}^t S(t-s)f(s, \varphi(s)) ds, \quad t \in \mathbb{R}.$$

Since $\varphi \in AA(X)$, we have that $\overline{\{\varphi(t) : t \in \mathbb{R}\}}$ is compact in X . By Theorem 2.10, there exists $q \in [1, p)$ such that $f(\cdot, \varphi(\cdot)) \in AS^q(X) \subset AS^1(X)$. Then, by Lemma 2.9, we conclude that F is well defined. Then for $\varphi_1, \varphi_2 \in AA(X)$ and $t \in \mathbb{R}$ we have:

$$\begin{aligned} \|F\varphi_1(t) - F\varphi_2(t)\| &\leq \int_{-\infty}^t \|S(t-s)\| \cdot \|f(s, \varphi_1(s)) - f(s, \varphi_2(s))\| ds \\ &\leq \int_0^\infty L(t-\tau) \|S(\tau)\| \cdot \|\varphi_1(t-\tau) - \varphi_2(t-\tau)\| d\tau \\ &\leq \|\varphi_1 - \varphi_2\|_\infty \int_0^\infty L(t-\tau) \phi(\tau) d\tau \\ &= \|\varphi_1 - \varphi_2\|_\infty \sum_{k=0}^\infty \int_k^{k+1} L(t-\tau) \phi(\tau) d\tau \\ &\leq \|\varphi_1 - \varphi_2\|_\infty \sum_{k=0}^\infty \phi(k) \int_k^{k+1} L(t-\tau) d\tau \\ &= \|\varphi_1 - \varphi_2\|_\infty \sum_{k=0}^\infty \phi(k) \int_{t-k}^{t-k-1} L(s) ds \\ &\leq \|\varphi_1 - \varphi_2\|_\infty \|L\|_{S^1} \sum_{k=0}^\infty \phi(k) \\ &\leq \|\varphi_1 - \varphi_2\|_\infty \|L\|_{S^1} \|\phi\|_1. \end{aligned}$$

This proves that F is a contraction, so by the Banach fixed point theorem there exists a unique $u \in AA(X)$, such that $Fu = u$, that is $u(t) = \int_{-\infty}^t S(t-s)f(s, u(s)) ds$. \square

We remark that in the case of $L(t) \equiv L$, by following the proof of previous theorem, one can get the same conclusion.

Our next result consider a Lipchitz condition with a function $L(t)$ whose norm does not need to be controlled, in contrast with our previous result. However, we have to add some extra conditions: boundedness of the resolvent family and integrability of $L(t)$.

Theorem 4.3. *Assume that A generates an integral resolvent family $\{S(t)\}_{t \geq 0}$ such that*

$$\|S(t)\| \leq \phi(t), \quad \text{for all } t \geq 0,$$

where $\phi \in L^1(\mathbb{R}_+) \cap L^\infty(\mathbb{R}_+)$ is nonincreasing. Suppose that

- (i) $f \in AS^p(\mathbb{R} \times X, X)$ with $p > 1$;
- (ii) there exists a non negative function $L \in AS^r(\mathbb{R}) \cap L^1(\mathbb{R})$ with $r \geq \max\{p, p/(p-1)\}$ such that for all $u, v \in X$ and $t \in \mathbb{R}$,

$$\|f(t, u) - f(t, v)\| \leq L(t)\|u - v\|;$$

Then equation (4.1) has a unique almost automorphic mild solution.

Proof. Define the operator F as in (4.3). Let φ_1, φ_2 be in $AA(X)$. We have:

$$\begin{aligned}
\|(F\varphi_1)(t) - (F\varphi_2)(t)\| &\leq \int_{-\infty}^t \|S(t-s)[f(s, \varphi_1(s)) - f(s, \varphi_2(s))]\| ds \\
&\leq \int_{-\infty}^t L(s) \|S(t-s)\| \|\varphi_1(s) - \varphi_2(s)\| ds \\
&\leq \|\varphi_1 - \varphi_2\|_{\infty} \int_{-\infty}^t L(s) \phi(t-s) ds \\
&= \|\varphi_1 - \varphi_2\|_{\infty} \int_0^{\infty} L(t-\tau) \phi(\tau) d\tau \\
&\leq \|\varphi_1 - \varphi_2\|_{\infty} \|\phi\|_{\infty} \int_0^{\infty} L(t-\tau) d\tau \\
&= \|\varphi_1 - \varphi_2\|_{\infty} \|\phi\|_{\infty} \int_{-\infty}^t L(s) ds.
\end{aligned}$$

In the same way

$$\begin{aligned}
\|(F^2\varphi_1)(t) - (F^2\varphi_2)(t)\| &\leq \int_{-\infty}^t \|S(t-s)[f(s, F\varphi_1(s)) - f(s, F\varphi_2(s))]\| ds \\
&\leq \int_{-\infty}^t L(s) \phi(t-s) \|F\varphi_1(s) - F\varphi_2(s)\| ds \\
&\leq \|\varphi_1 - \varphi_2\|_{\infty} \|\phi\|_{\infty}^2 \int_{-\infty}^t L(s) \int_{-\infty}^s L(\tau) d\tau ds.
\end{aligned}$$

In general we get

$$\begin{aligned}
\|(F^n\varphi_1)(t) - (F^n\varphi_2)(t)\| &\leq \|\varphi_1 - \varphi_2\|_{\infty} \frac{\|\phi\|_{\infty}^n}{(n-1)!} \left(\int_{-\infty}^t L(s) \left(\int_{-\infty}^s L(\tau) d\tau \right)^{n-1} ds \right) \\
&\leq \|\varphi_1 - \varphi_2\|_{\infty} \frac{\|\phi\|_{\infty}^n}{n!} \left(\int_{-\infty}^t L(s) ds \right)^n \\
&\leq \|\varphi_1 - \varphi_2\|_{\infty} \frac{(\|L\|_1 \|\phi\|_{\infty})^n}{n!}.
\end{aligned}$$

Hence, since $\frac{(\|L\|_1 \|\phi\|_{\infty})^n}{n!} < 1$ for n sufficiently large, by the contraction principle F has a unique fixed point $u \in AA(X)$. \square

Our next result, replaces the condition of integrability of $L(t)$ in the previous theorem by a less restrictive assumption (see (iii) below). The main idea is to define a new norm depending on a parameter (a weighted sup-norm) in $AA(X)$ in such way that the amount of the weight can be controlled, by means of the parameter, in the process of find a fixed point.

Theorem 4.4. *Assume that A generates an integral resolvent family $\{S(t)\}_{t \geq 0}$ such that*

$$\|S(t)\| \leq \phi(t), \text{ for all } t \geq 0,$$

where $\phi \in L^1(\mathbb{R}_+) \cap L^\infty(\mathbb{R}_+)$ is nonincreasing. Suppose that

(i) $f \in AS^p(\mathbb{R} \times X, X)$ with $p > 1$;

(ii) there exists a non negative function $L \in AS^r(\mathbb{R})$ with $r \geq \max\{p, p/(p-1)\}$ such that for all $u, v \in X$ and $t \in \mathbb{R}$,

$$\|f(t, u) - f(t, v)\| \leq L(t)\|u - v\|;$$

(iii) the integral $\int_{-\infty}^t L(s)ds$ exists for all $t \in \mathbb{R}$.

Then equation (4.1) has a unique almost automorphic mild solution.

Proof. Define a new norm $\|\varphi\| := \sup_{t \in \mathbb{R}} \{v(t)\|\varphi(t)\|\}$, where $v(t) := e^{-k \int_{-\infty}^t L(s)ds}$ and k is a fixed positive constant greater than $C := \|\phi\|_\infty$. Define the operator F as in (4.3). Let φ_1, φ_2 be in $AA(X)$, then we have

$$\begin{aligned} v(t)\|(F\varphi_1)(t) - (F\varphi_2)(t)\| &= v(t) \left\| \int_{-\infty}^t S(t-s)[f(s, \varphi_1(s)) - f(s, \varphi_2(s))]ds \right\| \\ &\leq C \int_{-\infty}^t v(t)L(s)\|\varphi_1(s) - \varphi_2(s)\|ds \\ &\leq C \int_{-\infty}^t v(t)v(s)^{-1}L(s)v(s)\|\varphi_1(s) - \varphi_2(s)\|ds \\ &\leq C\|\varphi_1 - \varphi_2\| \int_{-\infty}^t v(t)v(s)^{-1}L(s)ds \\ &= \frac{C}{k} \|\varphi_1 - \varphi_2\| \int_{-\infty}^t k e^{k \int_t^s L(\tau)d\tau} L(s)ds \\ &= \frac{C}{k} \|\varphi_1 - \varphi_2\| \int_{-\infty}^t \frac{d}{ds} \left(e^{k \int_t^s L(\tau)d\tau} \right) ds \\ &= \frac{C}{k} [1 - e^{-k \int_{-\infty}^t L(\tau)d\tau}] \|\varphi_1 - \varphi_2\| \\ &\leq \frac{C}{k} \|\varphi_1 - \varphi_2\|. \end{aligned}$$

Hence, since $C/k < 1$, F has a unique fixed point $u \in AA(X)$. \square

We finish this section with the following application, whose construction is inspired by Example 2.7.

Example 4.5. *Suppose that A is the generator of integral resolvent $\{S(t)\}_{t \geq 0}$ such that $\|S(t)\| \leq Me^{-\omega t}$ where $M, \omega > 0$. For concrete examples of operators satisfying this condition we refer to [26, Section 5]. For $\phi(t) = Me^{-\omega t}$, we observe that ϕ is nonincreasing and $\phi \in L^1(\mathbb{R}_+) \cap L^\infty(\mathbb{R}_+)$. Define*

$$l(t) := \begin{cases} \sin\left(\frac{1}{2+\cos n+\cos \pi n}\right), & t \in \left(n - \frac{\varepsilon}{2|n|}, n + \frac{\varepsilon}{2|n|}\right), n \in \mathbb{Z} \\ 0, & \text{otherwise,} \end{cases}$$

for some $0 < \varepsilon < \frac{1}{2}$. By Example 2.7, we conclude that $l \in AS^r(\mathbb{R})$ for all $r \in [1, \infty)$. Moreover,

$$\begin{aligned} \int_{-\infty}^{\infty} |l(t)| dt &\leq \sum_{n=-\infty}^{\infty} \int_{n-\frac{\varepsilon}{2^{|n|}}}^{n+\frac{\varepsilon}{2^{|n|}}} \left| \sin \left(\frac{1}{2 + \cos n + \cos \pi n} \right) \right| dt \\ &\leq \sum_{n=-\infty}^{\infty} \int_{n-\frac{\varepsilon}{2^{|n|}}}^{n+\frac{\varepsilon}{2^{|n|}}} dt \\ &= \sum_{n=-\infty}^{\infty} \frac{\varepsilon}{2^{|n|-1}} = 6\varepsilon. \end{aligned}$$

Therefore, $l \in AS^r(\mathbb{R}) \cap L^1(\mathbb{R})$. On $X = L^p([0, \pi])$, $p > 1$, define $f(t, u)(s) = l(t) \cos(u(s))$ for $u \in X$ and $s \in [0, \pi]$. Then, $f \in AS^p(\mathbb{R} \times X, X)$ and for $u, v \in X$, $t \in \mathbb{R}$ we have

$$\begin{aligned} \|f(t, u) - f(t, v)\|_p^p &= \int_0^\pi |l(t) \cos(u(s)) - l(t) \cos(v(s))|^p ds \\ &\leq |l(t)|^p \cdot \|u - v\|_p^p. \end{aligned}$$

We conclude from Theorem 4.3, with $L(t) = |l(t)|$, that the equation (4.1) has a unique almost automorphic mild solution. Note that, we obtain the same conclusion from Theorem 4.4. Finally, note that we can apply Theorem 4.2 under the condition $0 < \varepsilon < \frac{\omega}{2M}$.

Acknowledgement. We would like to thank the anonymous referee for his useful comments which lead to improve the paper.

REFERENCES

- [1] B. Basit, A.J. Pryde. *Asymptotic behavior of orbits of C_0 -semigroups and solutions of linear and semilinear abstract differential equations*, Russ. J. Math. Phys. **13** (1) (2006), 13–30.
- [2] D. Bugajewski, T. Diagana. *Almost automorphy of the convolution operator and applications to differential and functional differential equations*, Nonlinear Stud., **13** (2) (2006), 129–140.
- [3] B.D. Coleman, M.E. Gurtin. *Equipresence and constitutive equation for rigid heat conductors*, Z. Angew. Math. Phys., **18** (1967), 199–208.
- [4] C. Corduneanu. *Almost periodic solutions for infinite delay systems*, Spectral Theory of Differential Equations, I.W. Knowles and R.T. Davies (eds.), North Holland, 1981.
- [5] Ph. Clément, G. Da Prato. *Existence and regularity results for an integral equation with infinite delay in a Banach space*. Integral Equations Operator Theory, **11** (1988), 480–500.
- [6] Ph. Clément. *On abstract Volterra equations with completely positive kernels*. In: Infinite Dimensional Systems, Lecture Notes 1076, Springer, 1984.
- [7] Ph. Clément, P. Grisvard. *Sommes d'opérateurs linéaires et équations différentielles opérationnelles*. J. Math. Pures et Appl., **54** (1975), 305–388.
- [8] C. Cuevas, C. Lizama. *Almost automorphic solutions to integral equations on the line*. Semigroup Forum, **79** (3) (2009), 461–472.
- [9] E. Cuesta. *Asymptotic behaviour of the solutions of fractional integro-differential equations and some time discretizations*. Discrete Contin. Dyn. Syst. 2007, Dynamical Systems and Differential Equations. Proceedings of the 6th AIMS International Conference, suppl., 277–285.
- [10] T. Diagana. *Some remarks on some second-order hyperbolic differential equations*, Semigroup Forum **68** (2004), 357–364.
- [11] T. Diagana, G.M.N'Guérékata. *Stepanov-like almost automorphic functions and applications to some semilinear equations*, Appl. Anal. **86** (6) (2007) 723–733.

- [12] H. S. Ding, J. Liang, T. J. Xiao *Almost automorphic solutions to nonautonomous semilinear evolution equations in Banach spaces*, preprint.
- [13] H. S. Ding, J. Liang, T. J. Xiao *Almost automorphic solutions to abstract fractional differential equations*, Adv. Difference Equ. 2010, to appear.
- [14] J.A. Goldstein, G.M. N'Guérékata. *Almost automorphic solutions of semilinear evolution equations*, Proc. Amer. Math. Soc. **133** (8) (2005), 2401-2408.
- [15] R. Gorenflo, F. Mainardi. *On Mittag-Leffler-type functions in fractional evolution processes*, J. Comp. Appl. Math. **118** (2000), 283-299.
- [16] G. Gripenberg, S-O Londen, O. Staffans. *Volterra Integral and Functional Equations*. Encyclopedia of Mathematics and Applications, **34**, Cambridge University Press, Cambridge-New York, 1990.
- [17] Z. Hu, Z. Jin. *Stepanov-like pseudo-almost periodic mild solutions to perturbed nonautonomous evolution equations with infinite delay*. Nonlinear Anal. **71** (11) (2009), 5381-5391.
- [18] H. Lee, H. Alkahby. *Stepanov-like almost automorphic solutions of nonautonomous semilinear evolution equations with delay*. Nonlinear Anal. **69** (7) (2008), 2158-2166.
- [19] J. Liu, G.M. N'Guérékata, N. van Minh. *Almost automorphic solutions of second order evolution equations*, Appl. Anal. **84** (11) (2005), 1173-1184.
- [20] G.M. N'Guérékata. *Existence and uniqueness of almost automorphic mild solutions of some semilinear abstract differential equations*, Semigroup Forum **69** (2004), 80-86.
- [21] G.M. N'Guérékata. *Topics in Almost Automorphy*. Springer Verlag, New York, 2005.
- [22] G.M. N'Guerekata. *Almost Automorphic and Almost Periodic Functions in Abstract Spaces*. Kluwer Acad/Plenum, New York-Boston-Moscow-London, 2001.
- [23] G.M. N'Guerekata, A. Pankov. *Stepanov-like almost automorphic functions and monotone evolution equations*, Nonlinear Anal., **68** (2008) 2658-2667.
- [24] J. W. Nunziato. *On heat conduction in materials with memory*, Quart. Appl. Math., **29** (1971), 187-304.
- [25] A. Pankov. *Bounded and Almost Periodic Solutions of Nonlinear Operator Differential Equations*, Kluwer, Dordrecht, 1990.
- [26] J. Prüss. *Decay properties for the solutions of a partial differential equation with memory*. Arch. Math. (Basel) **92** (2) (2009), 158-173.
- [27] J. Prüss. *Evolutionary Integral Equations and Applications*. Monographs Math., **87**, Birkhäuser Verlag, 1993.

UNIVERSIDAD DE SANTIAGO DE CHILE, DEPARTAMENTO DE MATEMÁTICA, FACULTAD DE CIENCIAS,
CASILLA 307-CORREO 2, SANTIAGO-CHILE.

E-mail address: carlos.lizama@usach.cl

E-mail address: rodrigo.ponce@usach.cl